

Abduction Pillow

IF YOU EXPERIENCE ANY PAIN, SWELLING SENSATION CHANGES OR ANY UNUSUAL REACTIONS WHILE USING THIS PRODUCT, CONSULT YOUR MEDICAL PROFESSIONAL IMMEDIATELY. NEVER WEAR THE SUPPORT OVER AN OPEN WOUND.

Application

1. Detach the shoulder strap buckles and open front panel. Position elbow in the sling as far back as possible.
2. Place the shoulder strap over the opposite shoulder. Connect shoulder strap to the sling using the quick release buckles. For exercise and rehabilitation, detach the outside quick release buckle to increase the range of motion.
3. Secure the strap at the top of the sling. If the thumb strap is desired, attach the strap at the front of the sling.
4. Place the large pillow under the affected area with the 45° angle facing up. Attach the sling to the pillow, align the hook and loop straps. Wrap the waist strap around the waistline and the pillow and attach.
5. The pillow is to be worn 24/7, only coming off before bathing. Passive assisted movement of the shoulder will be allowed immediately after your surgery. The shoulder should be supported during times when you are without the pillow.

Use and Care

Hand wash with mild soap in lukewarm water. Rinse thoroughly. AIR DRY. Note: If not rinsed thoroughly, residual soap may cause skin irritation or deteriorate the material

Latex free. This product is intended for use on a single patient.

